

# CLINICAL TRIALS

---

## What clinical trials and scientific research supports Liquid BioCell™ Life's claims of healthy aging, active joints, and younger-looking skin?

- Acute and subchronic oral toxicity studies in rats of a hydrolyzed chicken sternal cartilage preparation. Schauss et al. 2007, *Food Chem Toxicol*, pp. 45:315-321.
- A randomized double blind clinical pilot trial evaluating the safety and efficacy of hydrolyzed collagen type II (Liquid BioCell™) in adults with osteoarthritis. Sheldon et al. 2003, Miami Research Associates.
- Boocock et al. Phase I dose escalation pharmacokinetic study in healthy volunteers of Resveratrol, a potential chemopreventive agent. *Cancer Epidemiol Biomarkers Prev.* 2007, Vols. 16(6): 1246-52.
- Articular cartilage: structure, injuries and review of management. Bhosale et al. 2008, *Brit Med Bull*, pp. 87: 77-95.
- Aggrecan, aging, and assembly in articular cartilage. Dudhia. 2005, *Cell Mol Life Sci*, pp. 62: 2241-2256.
- Effects of Oral Administration of Type II Collagen on Rheumatoid Arthritis. Trentham et al. 1993, *Science*, pp. 261: 1727-1730.
- Suppression of Type II Collagen-Induced Arthritis by the Intravenous Administration of Type II Collagen or Its Constituent Peptide. Englert et al. 1984, *Cell Immuno*, pp. 87: 357-365.
- Collagen hydrolysate for the treatment of osteoarthritis and other joint disorders: a review of the literature. Bello et al. 2006, *Curr Med Res Opin*, pp.22(11): 2221-2232.
- Hyaluronan synthesis and degradation in cartilage and bone. Bastow. 2008, *Cell Mol Life Sci*, pp. 65: 395-413.
- Stimulation of type II collagen biosynthesis and secretion in bovine chondrocytes cultured with degraded collagen. Oesser et al. 2003, *Cell Tissue Res*, pp. 399: 393-399.
- Detection of nitrotyrosine in aging and osteoarthritic cartilage: Correlation of oxidative damage with the presence of interleukin-1beta and with chondrocyte resistance to insulin-like growth factor 1. Loeser et al. 2002, *Arthritis Rheum* , pp. 46: 2349-2357.
- Oxidants, oxidative stress and the biology of aging. Finkel et al. 2000, *Nature*, pp. 408: 239-247.

- Potential involvement of oxidative stress in cartilage senescence and development of osteoarthritis: oxidative stress induces chondrocyte telomere instability and down-regulation of chondrocyte function. Yudoh et al. 2005, *Arthritis Res Ther*, pp. 7: R380-R391.
- Telomere erosion and senescence in human articular cartilage chondrocytes, Martin et al. 2001, *J Gerontol A Biol Sci Med Sci*, pp. 56: B172-B179.
- Loeser. Aging and osteoarthritis: the role of chondrocyte senescence and aging changes in the cartilage matrix. *Osteoarthritis and Cartilage*. 2009, Vols. 17:971-979.
- Hyaluronic acid in cutaneous intrinsic aging. Ghersetich et al. 1994, *Int J Dermatol*, pp. 33(2): 119-122.
- Stern et al. Hyaluronan catabolism: a new metabolic pathway. *Eur J Cell Biol*. 2004, Vols. 83(7): 317-325.
- Wiest et al. Native hyaluronic acid in dermatology-results of an expert meeting. *J Dtsch Dermatol Ges*. 2008, Vols. 6(3): 176-180.
- Collagen fragmentation promotes oxidative stress and elevates matrix metalloproteinase- 1 in fibroblasts in aged human skin. Fisher et al. 2009, *Am J Path*, pp. 174: 101-114.
- Chronic ultraviolet B irradiation causes loss of hyaluronic acid from mouse dermis because of down-regulation of hyaluronic acid synthesis. Dai et al. 2007, *Am J Path*, pp. 282: 1451-1561.
- Hyaluronic acid stimulates human fibroblast proliferation within a collagen matrix. Greco et al. 1998, *J Cell Physiol*, pp. 177(3): 465-473.
- Hyaluronan facilitates transforming growth factor-beta1-mediated fibroblastproliferation. Meran et al. 2008, *J Biol Chem*, pp. 283: 6530-6545.
- Aging: a theory based on free radical and radiation chemistry. Harman. 1956, *J Gerontol*, pp. 11(3): 298-300.
- Enzymatic and nonenzymatic antioxidants in epidermis and dermis of human skin. Shindo et al. 1994, *J Invest Dermatol*, pp. 102(1): 122-124.
- Intake of flavonoids and flavones and risk of coronary heart disease in male smokers. Hirvonen et al. 2001, *Epidemiology*, pp. 12: 62-67.
- In vivo exposure to ozone depletes vitamins C and E and induces lipid peroxidation in epidermal layers of murine skin. Thiele et al. 1997, *Free Radic Biol Med*, pp. 23(3): 385-91.
- Ascorbic acid specifically increases type I and type III procollagen messenger RNA levels in human skin fibroblast. Geesin et al. 1988, *J Invest Dermatol*, pp. 90(4): 420-424.28.
- Antioxidant activity of pomegranate juice and its relationship with phenolic composition and processing. Gil et al. 2000, *J Agric Food Chem*, pp. 48(10): 4581-4589.

- Studies on antioxidant activity of pomegranate (*Punica granatum*) peel extract using in vivo models. Chidambara et al. 2000, *J Agric Food Chem*, pp. 50(17): 4791-4795.
- Photochemopreventive effect of pomegranate fruit extract on UVA-mediated activation of cellular pathways in normal human epidermal keratinocytes. Syed et al. 2006, *Photochem Photobiol*, pp. 82(2): 398-405.
- Bog blueberry anthocyanins alleviate photoaging in ultraviolet-B irradiation- induced human dermal fibroblasts. Bae et al. 2009, *Mol Nutr Food Res*, pp.53(6): 726-738.
- Prevention of ultraviolet-B radiation damage by resveratrol in mouse skin is mediated via modulation in survivin. Aziz et al. 2005, *Phytochem Photobiol*, pp. 81(1): 25-31.
- Chemoprevention of skin cancer by grape constituent Resveratrol: relevance to human disease? Aziz et al. 2005, *FASEB J*, pp. 19(9): 1193-1195.34) A multitargeted agent for age-associated chronic diseases. Kuzhuvilil et al. 2008, *Cell Cycle*, p. 7:8.
- A low dose of dietary Resveratrol partially mimics caloric restriction and retards aging parameters in mice. Barger et al. 2008, *PLoS One*, p. 3(6): e2264.
- Schauss et al. 2012. Effect of the novel low molecular weight hydrolyzed chicken sternal cartilage extract, Liquid BioCell, on improving osteoarthritis-related symptoms: a randomized, double-blind, placebo-controlled trial. *J Agric Food Chem*. 60(16): 4096-101.
- Schwartz et al. 2012. Ingestion of Liquid BioCell, a novel hydrolyzed chicken sternal cartilage extract, enhanced blood microcirculation and reduced facial aging signs. *Clin Interv Aging*. 7: 267-273.